

Defining small sets from κ -cc families of subsets

I: Families of finite sets

Andrés Eduardo Caicedo, John D. Clemens,
Clinton Conley, and Benjamin D. Miller

December 4, 2009

Abstract

We show that if X is a set and $\mathcal{A} \subseteq [X]^n$ is non-empty and contains at most k pairwise disjoint elements, then a non-empty subset of X of cardinality strictly less than kn^2 is definable from \mathcal{A} . We show that this result is nearly best possible. We also study bounds on the size of intersecting families of ($\leq n$)-element subsets from which no smaller such family can be defined, and classify all families of this form for $n = 3$. Although guided by definability considerations, our arguments are combinatorial in nature.

Contents

1	Introduction	2
1.1	Application: A Glimm-Effros-style dichotomy	3
1.2	Acknowledgements	4
2	Intersecting families of finite sets	4
3	Minimal intersecting families	9
3.1	Examples	9
3.2	The size of minimal families	12
3.3	Minimal families of triples	16
4	Low intersecting families	24
A	Definability	27
	References	29

1 Introduction

Let X be a set, and let κ be a (possibly infinite) cardinal number. We say that a family $\mathcal{A} \subseteq \mathcal{P}(X)$ is an *antichain* iff

$$\forall A, B \in \mathcal{A} (A \neq B \Rightarrow A \cap B = \emptyset),$$

and we say that \mathcal{A} satisfies the κ -*chain condition*, or simply \mathcal{A} is κ -*cc*, iff it does not contain an antichain of cardinality κ . If $\kappa = 2$, the set \mathcal{A} is also called an *intersecting family*; if $\kappa = \aleph_0$, we say that \mathcal{A} is *finite-cc*. Notice that our notion of antichain is more restrictive than the alternative condition that elements of \mathcal{A} be \subseteq -incomparable.

Several results in diverse fields of mathematics depend on purely combinatorial principles. To name just two examples, Ramsey's theorem was established to prove a decidability result (and was rediscovered by Erdős and Szekeres to prove a result in geometry), and the study of automorphism groups of strongly regular graphs and Steiner triple systems played a role in the classification of the finite simple groups. It is often the case that the combinatorial principles behind results in other fields turn out to be interesting in their own right, and that their deeper study provides us with further applications of these principles.

A recent result of Clemens-Conley-Miller [2] in the theory of definable equivalence relations relies upon the ability to define finite sets in a canonical way from (possibly infinite) intersecting families of finite sets. Our aim in this paper is to investigate the purely combinatorial issues underlying the proof of this result, and in particular to study the minimum size of sets definable from such intersecting families. Extremal set theory, including the analysis of intersecting families, is a well known area of study within combinatorics; the twist we add is the consideration of definability conditions. This provides us with its own advantages, such as access to techniques from mathematical logic, and its own difficulties, such as the fact that probability arguments, Ramsey arguments, and non-constructive proofs do not yet seem applicable within our framework.

In §1.1, we briefly discuss the result from Clemens-Conley-Miller [2] leading to the combinatorial result that motivated this paper.

Definition 1.1. We denote by $\xi : \mathbb{Z}^+ \times \mathbb{Z}^+ \rightarrow \mathbb{Z}^+$ the map that assigns to each pair (k, n) the least l such that every non-empty $(k + 1)$ -cc family \mathcal{A} of n -element sets can be used to define a non-empty subset of $\bigcup \mathcal{A}$ of cardinality at most l .

In §2, we show that ξ is well defined, and provide bounds on its value. For this, we say that \mathcal{A} is a *low* family iff no proper subset of $\bigcup \mathcal{A}$ or of \mathcal{A} can be defined from \mathcal{A} ; our bounds come from the study of low families.

In §3, we introduce the concept of *minimal* intersecting families $\mathcal{A} \subseteq [X]^{\leq n}$. This is analogous to the concept of low but now rather than asking that no proper subset of X or \mathcal{A} is definable, we ask that no smaller intersecting family of $(\leq n)$ -element subsets of X (which need not even be a subfamily of \mathcal{A}) is definable. Let $\psi(n)$ be the size of the largest minimal family of $(\leq n)$ -element

sets. The arguments of §2 provide a super-exponential upper bound for ψ . We also find a super-exponential lower bound. In §3.3, we classify all minimal intersecting families of (≤ 3)-element sets. In particular, we show that if \mathcal{A} is such a family, then $|\mathcal{A}| \leq 10$, and show that there are exactly two non-isomorphic families for which this bound is attained. The classification also implies that $\xi(1, 3) = 7$, i.e., from any intersecting family of triples from a set X , one can define a subset of X of size at most 7, but not necessarily one of size 6.

We close the body of the paper in §4 by studying low intersecting families. We prove a lifting theorem that allows us to build larger low families from smaller ones. We also adapt an argument from §3.3 to show that if \mathcal{A} is a low family of 3-element sets, then $|\mathcal{A}| \leq 10$.

Finally, we formalize in Appendix A the concept of definability that our results deal with, and include the background material necessary to work with it.

In the sequel Caicedo-Clemens-Conley-Miller [1] we address similar problems for families of countable sets, both combinatorially, and in the context of descriptive set theory.

1.1 Application: A Glimm-Effros-style dichotomy

Recall that an equivalence relation E on a Polish space is said to be *countable* iff each of its equivalence classes $[x]_E$ is countable. A *reduction* from an equivalence relation E on X to another equivalence relation F on Y is a function $\pi : X \rightarrow Y$ with the property that $x_1 E x_2 \iff \pi(x_1) F \pi(x_2)$, for all $x_1, x_2 \in X$. The study of the Borel reducibility quasi-order (\leq_B) on the class of countable Borel equivalence relations on Polish spaces has played an important role in descriptive set theory over the last two decades.

A *partial transversal* of a countable Borel equivalence relation E is a set which intersects every E -class in at most one point. We denote by \mathcal{I}_E the σ -ideal consisting of those sets contained in the union of countably many Borel partial transversals of E . We say that E is *smooth* if \mathcal{I}_E trivializes, i.e., if $X \in \mathcal{I}_E$. By well-known results of classical descriptive set theory, such equivalence relations are \leq_B -minimal among all countable Borel equivalence relations on uncountable Polish spaces.

A remarkable result, which goes back to work of Glimm-Effros in operator algebras from the 1960s, is that there is a \leq_B -minimal non-smooth countable Borel equivalence relation on a Polish space. An example of such an equivalence relation is E_0 on $\{0, 1\}^{\mathbb{N}}$, which is given by

$$xE_0y \iff \exists n \in \mathbb{N} \forall m \geq n (x(m) = y(m)).$$

Speaking very roughly, this result is proved as follows. Given a countable Borel equivalence relation E , there is a natural attempt at recursively building a continuous injective reduction of E_0 into E , which essentially entails trying to build up copies of level-by-level approximations to E_0 within E . If this attempt fails to produce the desired reduction, then it necessarily provides a countable family of Borel partial transversals whose union is X , thus E is smooth.

Given both the central nature of the σ -ideal \mathcal{I}_E in this proof, as well as its alternative characterization as the σ -ideal generated by the Borel sets on which E is \leq_B -minimal, it is natural to ask for the extent to which \mathcal{I}_E determines the \leq_B -class of E . An answer to this question has been provided by Clemens-Conley-Miller [2], where it is showed that the existence of a Borel homomorphism from \mathcal{I}_E to \mathcal{I}_F is equivalent to the existence of a smooth-to-one Borel homomorphism from E to F (we refer the reader to Clemens-Conley-Miller [2] for the exact definitions). While the latter notion is strictly weaker than Borel reducibility, it does agree with Borel reducibility when restricted to several of the most important classes of countable Borel equivalence relations.

Lying at the heart of the Clemens-Conley-Miller [2] result is the fact that if E is a countable Borel equivalence relation on X , F is a countable Borel equivalence relation on Y , and $\phi : X \rightarrow Y$ is Borel, then either there is a Borel perturbation of ϕ which is a homomorphism from \mathcal{I}_E to \mathcal{I}_F , or else there is a continuous injective reduction π of E_0 into E with the property that the points of the form $\phi \circ \pi(x)$, for $x \in 2^{\mathbb{N}}$, are pairwise F -inequivalent. The proof resembles that of the Glimm-Effros theorem. Again, there is a natural attempt at recursively building the desired reduction of E_0 to E . This attempt fails if X is in the σ -ideal \mathcal{I}_ϕ generated by the Borel sets $B \subseteq X$ which have the property that for some finite set Γ of Borel automorphisms whose graphs are contained in E , the collection of sets of the form $\{[\phi(\gamma \cdot x)]_F : \gamma \in \Gamma\}$, for $x \in B$, is an intersecting family. In this case, one then obtains the desired Borel perturbation of ϕ by appealing to the fact that there is a Borel way of defining finite sets from intersecting families, which easily follows from the purely combinatorial result which we study in this paper.

1.2 Acknowledgements

The first author wants to thank the National Science Foundation for partial support through grant DMS-0801189.

2 Intersecting families of finite sets

Given a set X and a cardinal κ , let $[X]^\kappa = \{A \subseteq X : |A| = \kappa\}$, $[X]^{<\kappa} = \{A \subseteq X : |A| < \kappa\}$, and $[X]^{\leq \kappa} = \{A \subseteq X : |A| \leq \kappa\}$. We write $[X]^{<\mathbb{N}}$ for $[X]^{<\aleph_0}$. For each family $\mathcal{A} \subseteq \mathcal{P}(X)$ and $D \subseteq X$, let $\deg_{\mathcal{A}}(D)$ denote $|\{A \in \mathcal{A} : D \subseteq A\}|$. We often write $\deg_{\mathcal{A}}(x)$ instead of $\deg_{\mathcal{A}}(\{x\})$. An \mathcal{A} -*extension* of D (or simply an *extension* of D) is any $A \in \mathcal{A}$ with $D \subseteq A$.

For m a positive integer, and finite l , let

$$\mathcal{A}^{(m,l)} = \{D \in [X]^m : \deg_{\mathcal{A}}(D) > l\}.$$

Notice that $\mathcal{A}^{(m,l)}$ is definable from \mathcal{A} .

Example 2.1. Suppose that $A \cap B = \emptyset$ and $|A| \geq |B| = 3$. Let \mathcal{A} denote the family of sets of the form $\{a\} \cup (B \setminus \{b\})$, for $a \in A$ and $b \in B$. Then \mathcal{A} is

an intersecting family of 3-element sets and $\mathcal{A}^{(2,2)}$ is an intersecting family of 2-element sets.

Fact 2.2. *Suppose $\mathcal{A} \subseteq [X]^{<\mathbb{N}}$ is finite-cc. Then \mathcal{A} is $(k+1)$ -cc for some $k \in \mathbb{Z}^+$.*

Proof. Let $\mathcal{B} \subseteq \mathcal{A}$ be a maximal antichain, and let $k = |\bigcup \mathcal{B}|$. Let $\mathcal{C} \in [\mathcal{A}]^{k+2}$. Every non-empty $A \in \mathcal{C}$ contains an element of $\bigcup \mathcal{B}$ by maximality of \mathcal{B} , so at least two members of \mathcal{C} have an element in common, and \mathcal{C} is not an antichain. Of course, $k+2$ can be replaced with $k+1$ if $\emptyset \notin \mathcal{A}$. \square

Remark 2.3. Our assumption that the members of \mathcal{A} are finite cannot be relaxed in Fact 2.2. To see this, for each $k \in \mathbb{Z}^+$ let A_{k1}, \dots, A_{kk} be a partition of \mathbb{N} into k infinite sets such that if $k < l$ then $A_{ki} \cap A_{lj}$ is infinite for each i, j . Then $\mathcal{A} = \{A_{ki} : k \in \mathbb{Z}^+, 1 \leq i \leq k\}$ is finite-cc but not $(k+1)$ -cc for any $k \in \mathbb{Z}^+$.

For each non-empty $(k+1)$ -cc $\mathcal{A} \subseteq [X]^n$ and for each $m \leq n$, let $d_m = \sup_{D \in [X]^m} \deg_{\mathcal{A}}(D)$. In particular, $d_n = 1$. Let

$$\mathcal{A}_m = \mathcal{A}^{(m, (kn)d_{m+1})} = \{D \in [X]^m : \deg_{\mathcal{A}}(D) > (kn)d_{m+1}\}.$$

Lemma 2.4. *Suppose that $\mathcal{A} \subseteq [X]^n$ is $(k+1)$ -cc. Then for each $m < n$, \mathcal{A}_m is $(k+1)$ -cc (but possibly empty). Furthermore, if $\mathcal{A}_m \neq \emptyset$ and $\mathcal{A}_k = \emptyset$ for all $k > m$, then \mathcal{A}_m is definable from \mathcal{A} .*

Proof. Suppose, towards a contradiction, that there exist pairwise disjoint

$$D_0, \dots, D_k \in \mathcal{A}_m.$$

We inductively construct $A_0, \dots, A_k \in \mathcal{A}$ such that for each $i \leq k$:

1. A_i is an extension of D_i ,
2. for all $j > i$, $D_j \cap A_i = \emptyset$, and
3. for all $j < i$, $A_j \cap A_i = \emptyset$.

Suppose we have found A_0, \dots, A_{i-1} satisfying the above conditions. Notice that for any $x \in X \setminus D_i$, at most d_{m+1} extensions of D_i contain x , or else $\deg_{\mathcal{A}}(D_i \cup \{x\}) > d_{m+1}$. Consequently, no more than $(kn)d_{m+1}$ extensions of D_i can meet $\bigcup_{j < i} A_j \cup \bigcup_{j > i} D_j$, so there exists an extension A_i of D_i disjoint from $\bigcup_{j < i} A_j \cup \bigcup_{j > i} D_j$, completing this step of the construction. Of course, $\{A_0, \dots, A_k\}$ is an antichain of size $k+1$, contradicting the fact that \mathcal{A} is $(k+1)$ -cc.

It remains to show that if $m < n$ is largest such that $\mathcal{A}_m \neq \emptyset$, then \mathcal{A}_m is definable from \mathcal{A} . For this, argue by induction on $n-m$ that each d_i , $m < i \leq n$, is finite. In particular, d_{m+1} is finite, and the result follows. \square

Proposition 2.5. *Suppose that $\mathcal{A} \subseteq [X]^n$ is $(k+1)$ -cc. Then there exists a non-empty $(k+1)$ -cc family $\mathcal{A}' \subseteq [X]^{\leq n}$ definable from \mathcal{A} with $|\mathcal{A}'| \leq (kn)^n - kn + k$.*

Proof. The result is clear when $n = 1$ (with $\mathcal{A}' = \mathcal{A}$). We proceed by induction on n . If for some $m < n$, \mathcal{A}_m is non-empty, definable and $(k + 1)$ -cc, then the conclusion follows from the inductive hypothesis. Thus, by Lemma 2.4, we may assume that for all $m < n$, \mathcal{A}_m is empty. That is, for all $m < n$, $d_m \leq (kn)d_{m+1}$ and, in particular, $d_1 \leq (kn)^{n-1}$. Fix a maximal antichain $\{A_0, \dots, A_{k'-1}\}$ in \mathcal{A} , where $k' \leq k$. Every point in $\bigcup_{i < k'} A_i$ is contained in at most $(kn)^{n-1} - 1$ additional sets in \mathcal{A} . Since every set in \mathcal{A} intersects this maximal antichain, we have $|\mathcal{A}| \leq (k'n)((kn)^{n-1} - 1) + k'$, thus $|\mathcal{A}| \leq (kn)^n - kn + k$. \square

In particular, Proposition 2.5 implies the (crude) bound

$$\xi(k, n) \leq n((kn)^n - kn + k),$$

where ξ is as in Definition 1.1. We now give an improved bound as a function of k , n , and d_1 .

Proposition 2.6. *Suppose that $\mathcal{A} \subseteq [X]^n$ is $(k + 1)$ -cc with d_1 finite, and let $Y = \{x \in X : \deg_{\mathcal{A}}(x) = d_1\}$. Then*

$$|Y| \leq k(n^2 - (n/d_1)(n - 1)).$$

Proof. Simply observe that

$$\begin{aligned} \frac{d_1|Y|}{n} &\leq \frac{1}{n} \sum_{x \in X} \deg_{\mathcal{A}}(x) \\ &= |\mathcal{A}| \\ &\leq k(1 + n(d_1 - 1)), \end{aligned}$$

thus $|Y| \leq k(n^2 - (n/d_1)(n - 1))$. \square

As a corollary, we see that $\xi(k, n) \leq kn^2 - 1$, which is a quantitative improvement of the observation of Clemens-Conley-Miller [2] that gave rise to this paper:

Theorem 2.7. *Suppose that k and n are positive integers.*

1. *Suppose that $\mathcal{A} \subseteq \mathcal{P}(X)$ is finite-cc, $\mathcal{A} \cap [X]^n \neq \emptyset$, and $\mathcal{A} \cap [X]^n$ admits a maximal antichain of size k . Then a non-empty subset of X of cardinality at most $kn^3 - 1$ is definable from \mathcal{A} .*
2. *If $\mathcal{A} \subseteq [X]^n$ is $(k + 1)$ -cc, then a non-empty subset of X of cardinality at most $kn^2 - 1$ is definable from \mathcal{A} .*

Proof. We prove (1); the argument for (2) is identical. By replacing \mathcal{A} with $\mathcal{A} \cap [X]^n$, we can assume that $\mathcal{A} \subseteq [X]^n$. By Fact 2.2, \mathcal{A} is $(kn + 1)$ -cc. By Proposition 2.5, we can assume that d_1 is finite, and Proposition 2.6 then gives the desired result. \square

Corollary 2.8. *Suppose that $\mathcal{A} \subseteq [X]^{<\mathbb{N}}$ and that there exist non-empty (not necessarily definable) sets $\mathcal{A}_0, \mathcal{A}_1$ such that $\mathcal{A} = \mathcal{A}_0 \cup \mathcal{A}_1$ and for all $A_0 \in \mathcal{A}_0$ and $A_1 \in \mathcal{A}_1$, $A_0 \cap A_1 \neq \emptyset$. Then there is a non-empty, finite subset of X definable from \mathcal{A} .*

Proof. Notice that the assumption ensures that $\emptyset \notin \mathcal{A}$. In light of Theorem 2.7.1, it suffices to show that \mathcal{A} is finite-cc. For this, simply note that any antichain \mathcal{C} is contained in \mathcal{A}_i for some $i \in \{0, 1\}$. Since every element of \mathcal{C} intersects every $A \in \mathcal{A}_{1-i}$, we see that $|\mathcal{C}| \leq |A|$ for any such A . \square

Remark 2.9. The proof of Corollary 2.8 in fact shows that we can find a definable set of size at most

$$\left(\max_{i < 2} \min_{A \in \mathcal{A}_i} |A|\right) \left(\min_{A \in \mathcal{A}} |A|\right)^2 - 1.$$

We now show that the bound of $kn^2 - 1$ from Theorem 2.7.2 is best possible, in the sense that we cannot replace it with ϵkn^2 for any $0 < \epsilon < 1$, using some well-known examples from the study of combinatorial designs.

Recall that the notion of low was introduced in Definition A.7. Notice that $\bigcup \mathcal{A}$ is definable from \mathcal{A} , so if \mathcal{A} is low, then $X = \bigcup \mathcal{A}$. We say that \mathcal{A} has *degree* k iff $\forall x \in X$ ($\deg_{\mathcal{A}}(x) = k$). If the degree of \mathcal{A} is defined (for example, if \mathcal{A} is low), then \mathcal{A} is called *regular*.

Theorem 2.10. *Suppose that k and n are positive integers.*

1. *There is a set X of cardinality $k(n^2/2 + n/2)$ and a low $(k+1)$ -cc family $\mathcal{A} \subseteq [X]^n$ of degree 2.*
2. *If $n-1$ is a power of a prime, then there is a set X of cardinality $k(n^2 - n + 1)$ and a low $(k+1)$ -cc family $\mathcal{A} \subseteq [X]^n$ of degree n .*

Proof. We will handle only the case that $k = 1$, as the general case follows by taking the disjoint union of k copies of the examples we describe. To see (1), let $I = \{0, 1, \dots, n\}$ and $X = [I]^2$. Then

$$|X| = n^2/2 + n/2.$$

For each $i \in I$, define

$$A_i = \{x \in X : i \in x\},$$

so $|A_i| = n$ and

$$\mathcal{A} = \{A_i : i \in I\} \subseteq [X]^n$$

is an intersecting family of degree 2 since if $x = \{i, j\} \in X$, then $A_i \cap A_j = \{x\}$ and $x \in A_k \iff k \in x$. To see that \mathcal{A} is low, define

$$G = \{g \in S_X : \exists \sigma \in S_I \forall \{i, j\} \in X (g \cdot \{i, j\} = \{\sigma \cdot i, \sigma \cdot j\})\},$$

and observe that $G \leq \text{Aut}(\mathcal{A})$ and G acts transitively on X and \mathcal{A} .

To see (2), set $q = p^k = n - 1$. Let \mathbb{F}_q be the field of size q , and let V be a vector space of dimension 3 over \mathbb{F}_q .

Let

$$X = \{W \leq V : \dim W = 1\},$$

so

$$|X| = (q^3 - 1)/(q - 1) = q^2 + q + 1 = n^2 - n + 1,$$

since two non-zero vectors \mathbf{x} and \mathbf{y} on V determine the same line W iff there is $k \in \mathbb{F}_q \setminus \{0\}$ such that $\mathbf{x} = k\mathbf{y}$.

For $Y \leq V$ of dimension 2, let $\bar{Y} = \{W \leq Y : \dim W = 1\}$ and set

$$\mathcal{A} = \{\bar{Y} : Y \leq V, \dim Y = 2\}.$$

Notice that

$$|\bar{Y}| = \frac{q^2 - 1}{q - 1} = q + 1 = n.$$

We can endow V with a formal inner product by fixing an arbitrary basis B of V over \mathbb{F}_q and setting $\mathbf{x} \cdot \mathbf{y} = \sum_i x_i y_i$ where $\langle x_1, x_2, x_3 \rangle$ and $\langle y_1, y_2, y_3 \rangle$ are the coordinates of \mathbf{x} and \mathbf{y} , respectively, with respect to the basis B . The map that sends W to its orthogonal complement is a one-to-one correspondence between spaces W of dimension 1 and spaces Y of dimension 2, so

$$|\mathcal{A}| = |X|.$$

Clearly, \mathcal{A} is intersecting. Given W of dimension 1 and Y of dimension 2, let Y_\perp be the orthogonal complement of Y and let W^\perp be the orthogonal complement of W . Then $W \in \bar{Y}$ iff $Y = (W^\perp)_\perp$, so

$$\deg_{\mathcal{A}}(W) = |\overline{W^\perp}| = n.$$

(For $k = 1$, what we are describing should remind the reader of a well known Steiner $S(2, p - 1, p^2 + p + 1)$ system, see van Lint-Wilson [6].) To see that \mathcal{A} is low, define

$$G = \{g \in S_X : \exists T \in \text{GL}_3(\mathbb{F}_q) \forall W \in X (g \cdot W = TW)\},$$

and observe that $G \leq \text{Aut}(\mathcal{A})$ and G acts transitively on X and \mathcal{A} . □

Example 2.11. When $n = 3$ and $k = 1$, Theorem 2.10.2 provides us with a low intersecting family of triples with $|X| = 7$, the *Fano plane*. This can also be described by letting $X = \mathbb{Z}/7\mathbb{Z}$ and taking as \mathcal{A} the family of translations of $\{0, 1, 3\}$:

$$\mathcal{A} = \{013, 124, 235, 346, 450, 561, 602\}.$$

This shows that $\xi(1, 3) \geq 7$. We in fact have equality, see Proposition 3.41.

3 Minimal intersecting families

For the remainder of the paper, we confine our attention to intersecting families. In the previous section we found bounds for the smallest size of a subset of X definable from \mathcal{A} , where $\mathcal{A} \subseteq [X]^n$ is intersecting. This was done in two stages; in the first, we found a finite intersecting $\mathcal{A}' \subseteq [X]^{\leq n}$ definable from \mathcal{A} . In the second, we started from such a finite \mathcal{A}' and found bounds on the size of subsets of X definable from \mathcal{A}' (and therefore, from \mathcal{A}). Here we address the question of how small we can find \mathcal{A}' itself.

Definition 3.1. An intersecting family $\mathcal{A} \subseteq [X]^{\leq n}$ is *n-minimal* (or, simply, *minimal* if n is clear from context) iff there is no intersecting $\mathcal{B} \subseteq [X]^{\leq n}$ definable from \mathcal{A} with $|\mathcal{B}| < |\mathcal{A}|$.

Remark 3.2. Given a set X and intersecting families $\mathcal{A}_0, \mathcal{A}_1 \subseteq \mathcal{P}(X)$, the relation “ \mathcal{A}_0 is definable from \mathcal{A}_1 ” is a quasi-order. It is not a partial order since, for example, if $X = \{0, 1, 2\}$, then $[X]^2$ and $[X]^3$ can both be defined from each other.

If \mathcal{A} is minimal, we may assume $\bigcup \mathcal{A} = X$, so we adopt this convention (often without comment) in what follows; this is a minor technicality intended to avoid situations like the following: We could have a set X of size at least 10, let $Y \subset X$ have size 5, and let $\mathcal{A} = [Y]^3$. Then \mathcal{A} is 3-minimal and of size 10. We have that $\bigcup \mathcal{A} = Y$ and $[X \setminus Y]^3$ is definable from \mathcal{A} ; moreover, if $|X| = 10$, then $[X \setminus Y]^3$ is intersecting and of size 10 as well. Obviously, $X \setminus Y$ has nothing to do with the combinatorics of \mathcal{A} , so we prefer to discard it rather than having to pay attention to its size.

Notice that if $\mathcal{A} \subseteq [X]^{\leq n}$ is minimal, then there is $m \leq n$ such that $\mathcal{A} \subseteq [X]^m$.

Definition 3.3. Let $\psi(n)$ be the largest possible size of an n -minimal intersecting family of sets. A family $\mathcal{A} \subseteq [X]^{\leq n}$ is said to be *n-large* (or, simply, *large* if n is clear from context) if it is intersecting, minimal, and has size $\psi(n)$.

Notice that $\psi(n)$ is finite for all n . In fact, $\psi(n) \leq n^n - n + 1$, by Proposition 2.5.

3.1 Examples

Example 3.4. Let $X = \{1, \dots, 2n - 1\}$ and $\mathcal{A} = [X]^n$. Then \mathcal{A} is intersecting, low and minimal. Hence

$$\binom{2n - 1}{n} \leq \psi(n).$$

It follows that $\psi(2) \geq 3$ and $\psi(3) \geq 10$. It is easy to see that, in fact, $\psi(2) = 3$.

Example 3.5. Besides the family of 3-element subsets of a set of size 5, there is another example of a minimal family of triples of size 10: Let $X = \{a, b, c, d, e, f\}$ and consider the family

$$\mathcal{A} = \{abc, abf, ace, ade, bcd, bde, bef, cdf, cef\}.$$

To see that \mathcal{A} is minimal, first we show that it is low. For this, it suffices to consider the subgroup of $\text{Aut}(\mathcal{A})$ generated by the permutations π_1 and π_2 , where

- $\pi_1(a) = b, \pi_1(b) = e, \pi_1(c) = d, \pi_1(d) = a, \pi_1(e) = c$ and $\pi_1(f) = f$, and
- $\pi_2(a) = c, \pi_2(b) = d, \pi_2(c) = f, \pi_2(d) = a, \pi_2(e) = e$ and $\pi_2(f) = b$.

That is, when presented in cycle notation,

- $\pi_1 = (abecd)(f)$, and
- $\pi_2 = (acfbd)(e)$.

But then it follows that \mathcal{A} is minimal: Suppose that $\mathcal{B} \subseteq [X]^{\leq 3}$ is intersecting and definable from \mathcal{A} . If $\mathcal{B} \cap [X]^{\leq 2} \neq \emptyset$, then from $\mathcal{B} \cap [X]^{\leq 2}$ (and, therefore, from \mathcal{A}) we can define a non-empty subset of X of size at most 3, contradicting that \mathcal{A} is low. Thus, $\mathcal{B} \subseteq [X]^3$. If $\mathcal{B} \cap \mathcal{A} \neq \emptyset$ then $\mathcal{B} \cap \mathcal{A} = \mathcal{A}$, since \mathcal{A} is low, but then $|\mathcal{B}| \geq |\mathcal{A}|$. So we may assume that $\mathcal{B} \cap \mathcal{A} = \emptyset$. But then, again considering $\langle \pi_1, \pi_2 \rangle$, $\mathcal{B} \supseteq [X]^3 \setminus \mathcal{A}$, so $|\mathcal{B}| = 10 = |\mathcal{A}|$. (Notice that $[X]^3 \setminus \mathcal{A}$ is intersecting and definable from \mathcal{A} , so this last possibility occurs.)

Notice that under the action of S_X on $\mathcal{P}(\mathcal{P}(X))$, the permutation π given by

$$\pi(a) = d, \pi(b) = e, \pi(c) = f, \pi(d) = a, \pi(e) = b, \pi(f) = c$$

(i.e., $\pi = (ad)(be)(cf)$) exchanges \mathcal{A} and \mathcal{B} , showing that $\hat{\mathcal{A}}_X$ and $\hat{\mathcal{B}}_X$ are isomorphic.

Example 3.6. The previous example is a particular case of a more general construction of intersecting families, which is worth considering in some detail. Fix n and let $X = \{0, 1, \dots, 2n - 1\}$. Let $\mathcal{A} \subset [X]^n$ be such that for any $A \in [X]^n$ either A or $X \setminus A$ is not in \mathcal{A} . Then \mathcal{A} is intersecting and

$$|\mathcal{A}| \leq \frac{1}{2} \binom{2n}{n} = \binom{2n-1}{n}.$$

If we want \mathcal{A} to be of maximal size and minimal, our choice as to which one of A and $X \setminus A$ belongs to \mathcal{A} needs to be sufficiently uniform. For example, let $n = 3$. For $A \subseteq X$ let $\sum A = \sum_{x \in A} x$. Since $\sum X = 15$ is odd, we can define an \mathcal{A} of maximal size by letting

$$\mathcal{A} = \{A \in [X]^3 : \sum A \text{ is even}\}.$$

However, this \mathcal{A} is neither minimal nor low, since $\deg_{\mathcal{A}}(0) = 4$ while $\deg_{\mathcal{A}}(1) = 6$.

We can use a similar idea to produce (at least for $n = 3$) families that are actually minimal. Once again, let n be arbitrary and notice that for any $A \subseteq X$, $\sum A + \sum(X \setminus A) = n(2n - 1)$, so $\sum A + \sum(X \setminus A) \equiv 0 \pmod{n}$. If n is odd, we can then fix a set S such that for all $i \neq 0$, exactly one of $i, -i$ is in $S \pmod{n}$.

Then we would want to take $\mathcal{A} = \{A \in [X]^n : \sum A \in S \pmod{n}\}$. However, this does not take care of the case when $\sum A \equiv \sum(X \setminus A) \equiv 0 \pmod{n}$, but in that case, since $n(2n-1)$ is an odd multiple of n , exactly one of $\sum A, \sum(X \setminus A)$ is an odd multiple of n . So we just choose one of these options and put the corresponding set into \mathcal{A} .

Applying this idea to the case $n = 3$ we obtain the following families:

- $\{A \in [X]^3 : \sum A \equiv 1 \pmod{3} \text{ or } \sum A \equiv 3 \pmod{6}\}$.
- $\{A \in [X]^3 : \sum A \equiv 2 \pmod{3} \text{ or } \sum A \equiv 3 \pmod{6}\}$.
- $\{A \in [X]^3 : \sum A \equiv 1 \pmod{3} \text{ or } \sum A \equiv 0 \pmod{6}\}$.
- $\{A \in [X]^3 : \sum A \equiv 2 \pmod{3} \text{ or } \sum A \equiv 0 \pmod{6}\}$.

It is easy to verify that these four families are all isomorphic to the family from Example 3.5.

Leaving aside issues of minimality, this construction obviously generalizes to larger odd values of n . However, it is not possible to do something like this and obtain a minimal family for all even values of n :

Lemma 3.7. *Let $\mathcal{A} \subseteq [X]^{\leq n}$ be minimal. Assume that $|\mathcal{A}| \geq \binom{2n-1}{n}$ and $|X| \leq 4n-3$. Then \mathcal{A} is low. If $|\mathcal{A}| > \binom{2n-1}{n}$, the same holds for $|X| \leq 4n-1$.*

Proof. Since \mathcal{A} is minimal, by Lemma A.12, that \mathcal{A} is not low is equivalent to there being a non-empty proper subset of X definable from \mathcal{A} . But, if there were such a set A , either A or its complement would have size at most $2n-2$ (or at most $2n-1$ if $|\mathcal{A}| > \binom{2n-1}{n}$). Without loss of generality, assume it is A . Then the family of n -element subsets of A would be intersecting and of size strictly smaller than \mathcal{A} , contradicting minimality. \square

For example, it follows that for $n = 4$ we would need a family of 35 4-element subsets of a set of size 8 where each element has the same degree d , but this would imply that $8d = 4 \times 35$, a contradiction. (On the other hand, $\binom{2n-1}{n}$ is always even for odd $n > 1$.)

As for the question of whether \mathcal{A} is regular, we can say the following: For definiteness, let $n = 2k-1$ and set $S = \{1, \dots, k-1\}$ and

$$\mathcal{A} = \{A \in [\{0, \dots, 2n-1\}]^n : \sum A \in S \pmod{n} \text{ or } \sum A \equiv n \pmod{2n}\}.$$

The map $i \mapsto i+2 \pmod{2n}$ is in $\text{Aut}(\mathcal{A})$, so even numbers are indistinguishable, and so are odd numbers. The map $i \mapsto n-i$ is an isomorphism between \hat{A}_X and $\widehat{A^c}_X$ that exchanges the roles of even and odd numbers. The map $i \mapsto i+1$ sends \mathcal{A} to $\bar{\mathcal{A}} = \{A : \sum A \in S \pmod{n} \text{ or } \sum A \equiv 0 \pmod{2n}\}$, again exchanging the roles of even and odd numbers. It follows that \mathcal{A} is regular iff $|\{A \in \mathcal{B} : 0 \in A\}| = |\{A \in \mathcal{B} : 1 \in A\}|$, where

$$\mathcal{B} = \{A \in [\{0, \dots, 2n-1\}]^n : \sum A \equiv n \pmod{2n}\}.$$

However, examination of these families for small values of n has failed to provide examples of regular collections \mathcal{A} other than for $n = 3$.

3.2 The size of minimal families

Since minimal intersecting families are necessarily finite, in the current subsection we restrict our attention to finite intersecting families. Given a finite, intersecting $\mathcal{A} \subseteq [X]^{\leq n}$, recall that

$$d_1 = \max\{\deg_{\mathcal{A}}(x) : x \in X\}$$

and set

$$S_1 := \{x : \deg_{\mathcal{A}}(x) = d_1\}.$$

Question 3.8. *Assume $\mathcal{A} \subseteq [X]^{\leq n}$ is large. Does it follow that $\mathcal{A} \subseteq [X]^n$? Equivalently, is ψ (strictly) increasing?*

Fact 3.9. *Assume $n > 1$ and $\psi(n) < \psi(n+1)$. Let $\mathcal{A} \subseteq [X]^{n+1}$ be a minimal intersecting family with $|\mathcal{A}| > \psi(n)$. Then $\deg_{\mathcal{A}}(x) > 1$ for all $x \in X$.*

Proof. Otherwise, the collection \mathcal{B} of sets $Y \subset X$ of size n such that there is an $x \in X$ with $\deg_{\mathcal{A}}(x) = 1$ and $Y \cup \{x\} \in \mathcal{A}$ is intersecting and definable from \mathcal{A} . From it, an intersecting family \mathcal{C} of n -element subsets of X of size at most $\psi(n)$ can be defined, by definition of ψ . But since $\psi(n) < |\mathcal{A}|$, this contradicts minimality of \mathcal{A} . \square

Note that if \mathcal{A} is minimal, then $m_1 := |A \cap S_1|$ is independent of $A \in \mathcal{A}$.

Lemma 3.10. *Let $\mathcal{A} \subseteq [X]^{\leq n}$ be minimal, $|\mathcal{A}| \geq \binom{2n-1}{n}$. Then $|S_1| \geq 2n-1$, with equality only if $|\mathcal{A}| = \binom{2n-1}{n}$. If $n > 1$ then $m_1 > 1$.*

Proof. If $|S_1| < 2n-1$ then $[S_1]^n$ is intersecting and of size strictly smaller than $|\mathcal{A}|$, contradicting the minimality of \mathcal{A} . If $|\mathcal{A}| > \binom{2n-1}{n}$, then in fact $|S_1| \geq 2n$ by the same argument.

Assume $m_1 = 1$. Fix $x \in S_1$. There are at least $2n-2$ other points in S_1 , so at least $(2n-2)d_1$ sets in \mathcal{A} not containing x . Fix $A \in \mathcal{A}$ with $x \in A$. If $n > 1$, these $(2n-2)d_1$ sets all meet $A \setminus \{x\}$, so one of the points of $A \setminus \{x\}$ has degree at least $(2n-2)d_1/(n-1) > d_1$, contradiction. \square

Fact 3.11. *Let \mathcal{A} be n -minimal, and let $d_1 = d_{1,1} > \dots > d_{1,k}$ be the degrees of elements of X . Set $S_{1,i} := \{x : \deg_{\mathcal{A}}(x) = d_{1,i}\}$.*

1. *Every $A \in \mathcal{A}$ has elements of each degree (so $k \leq n$).*
2. *$m_i := |A \cap S_{1,i}|$ is independent of $A \in \mathcal{A}$.*
3. $\frac{d_{1,i}|S_{1,i}|}{m_i} = |\mathcal{A}|$.

Proof. The first two assertions are clear from the minimality of \mathcal{A} . The last one follows from a double counting argument:

$$\begin{aligned} d_{1,i}|S_{1,i}| &= \sum_{x \in S_{1,i}} d_{1,i} = \sum_{x \in X} \sum_{A \in \mathcal{A}} \chi_{S_{1,i}}(x) \chi_A(x) \\ &= \sum_A \sum_x \chi_{S_{1,i}}(x) \chi_A(x) = \sum_A |S_{1,i} \cap A| \\ &= m_i |\mathcal{A}|, \end{aligned}$$

and we are done. \square

Fact 3.12. *Assume that $\psi(n-1) < \psi(n)$. Let $\mathcal{A} \subseteq [X]^{\leq n}$ be a minimal intersecting family with $|\mathcal{A}| > \psi(n-1)$. Let*

$$S_{n-1} := \{D \in [X]^{n-1} : \deg_{\mathcal{A}}(D) = d_{n-1}\}.$$

Then $d_{n-1} \leq n$ and if $n > 2$ and $d_{n-1} = n$, then S_{n-1} is 3-cc.

Proof. That $d_{n-1} \leq n$ follows from Lemma 2.4 and the fact that $|\mathcal{A}| > \psi(n-1)$.

Assume now that $d_{n-1} = n > 2$ and that $D_1, D_2, D_3 \in S_{n-1}$ are pairwise disjoint. We can then find x_1, x_3 such that $D_i \cup \{x_i\} \in \mathcal{A}$ and $x_i \notin D_2$, $i = 1, 3$. Since $d_{n-1} = n = |D_3 \cup \{x_3\}|$ and any extension of D_2 meets $D_3 \cup \{x_3\}$, it follows that, for any $a \in D_3$, $D_2 \cup \{a\} \in \mathcal{A}$ and therefore it meets $D_1 \cup \{x_1\}$, so $x_1 = a$. But this contradicts the fact that $|D_3| = n-1 > 1$. \square

At this point, it is worth mentioning a reformulation of Lemma 2.4 for finite intersecting families \mathcal{A} . We first generalize a notion introduced earlier.

Definition 3.13. Let $\mathcal{A} \subseteq [X]^{\leq n}$. For $m \leq n$ let

$$S_m = \{D \in [X]^m : \deg_{\mathcal{A}}(D) = d_m\}.$$

Since for all $m < n$, $d_m \leq |\mathcal{A}|$, every S_m is definable from \mathcal{A} . With the nuisance of definability out of the way, Lemma 2.4 can be stated much more simply:

Lemma 3.14. *Let $\mathcal{A} \subseteq [X]^n$ be intersecting and finite. For all $m < n$ at least one of the following holds:*

1. S_m is intersecting.

2. $d_m \leq nd_{m+1}$. \square

The above suggests that the larger \mathcal{A} is, the smaller X seems required to be, if we want \mathcal{A} to be minimal. This intuition is not completely accurate, as the remainder of this subsection shows.

Question 3.15. *Is a large $\mathcal{A} \subseteq [X]^{\leq n}$ necessarily low?*

Definition 3.16. A family $\mathcal{A} \subseteq [X]^n$ is *strongly minimal* or *sminimal* iff $\text{Aut}(\mathcal{A})$ acts transitively on \mathcal{A} and on $[X]^m$ for all $m < n$.

Question 3.17. *If \mathcal{A} is large, is it sminimal?*

Theorem 3.18. *Suppose $\mathcal{A} \subseteq [X]^n$ is intersecting and $|\mathcal{A}| > \psi(n-1)$. Then the following are equivalent:*

1. \mathcal{A} is minimal.

2. For all $A \in [X]^{\leq n}$ either $A \notin \mathcal{A}$ and there is $\sigma \in \text{Aut}(\mathcal{A})$ such that $A \cap \sigma \cdot A = \emptyset$, or else $A \in [X]^n$ and $|\text{Aut}(\mathcal{A}) \cdot A| \geq |\mathcal{A}|$.

Proof. (1 \Rightarrow 2) Suppose \mathcal{A} is minimal so, in particular, it is finite. Let $A \in [X]^{\leq n}$. Let $\mathcal{B} = \mathcal{B}_A := \text{Aut}(\mathcal{A}) \cdot A$ be the closure of A under $\text{Aut}(\mathcal{A})$. By Lemma A.12, \mathcal{B} is definable from \mathcal{A} and is contained in every subset of $\mathcal{P}(X) \cup X$ definable from \mathcal{A} that contains A as an element. By minimality of \mathcal{A} , either \mathcal{B} is not intersecting, which means that for some $\sigma \in \text{Aut}(\mathcal{A})$, $A \cap \sigma \cdot A = \emptyset$, or else $|\mathcal{B}| = |\text{Aut}(\mathcal{A}) \cdot A| \geq |\mathcal{A}|$. In the first case, $A \notin \mathcal{A}$. Otherwise, $\mathcal{B} \subseteq \mathcal{A}$ (since \mathcal{A} is clearly definable from \mathcal{A} and contains A as an element), but \mathcal{B} is not intersecting, a contradiction. In the second case, it follows that $|A| = n$. Otherwise, there is an intersecting family \mathcal{C} of $\leq (n-1)$ -element sets with $|\mathcal{C}| \leq \psi(n-1) < |\mathcal{A}|$ definable from \mathcal{B} (and therefore from \mathcal{A}), contradicting the minimality of \mathcal{A} .

(2 \Rightarrow 1) Let \mathcal{A} satisfy the second condition, and suppose \mathcal{A} is not minimal. Let \mathcal{B} be an intersecting family of $(\leq n)$ -element sets definable from \mathcal{A} with $|\mathcal{B}| < |\mathcal{A}|$. For any $A \in \mathcal{B}$, $\mathcal{B}_A \subseteq \mathcal{B}$, so \mathcal{B}_A is intersecting. It follows that $|A| = n$ and $|\mathcal{B}_A| \geq |\mathcal{A}|$, contradicting that $|\mathcal{B}| < |\mathcal{A}|$. \square

A similar argument gives the following sufficient condition for minimality:

Lemma 3.19. *Let $\mathcal{A} \subseteq [X]^n$ be intersecting. Suppose that $\text{Aut}(\mathcal{A})$ is transitive on \mathcal{A} and for all $B \in [X]^{\leq n} \setminus \mathcal{A}$ there is a $\sigma \in \text{Aut}(\mathcal{A})$ such that $B \cap \sigma \cdot B = \emptyset$. Then \mathcal{A} is minimal.* \square

Theorem 3.20. *For all even $n > 6$, $\psi(n) > (n/2)^{n/2}$ and there is a minimal intersecting $\mathcal{A} \subseteq [X]^n$ with $|X| \geq n^2/4$. In fact, for all rational $p \in (0, 1)$, for all but finitely many n such that pn is an integer, $\psi(n) > (pn)^{(1-p)n+1}$.*

Proof. Assume $n = 2k$ is even. Let $X = \bigsqcup_{i=0}^k X_i$ be the disjoint union of sets X_i with $|X_i| = k$. For $I \subseteq \{0, \dots, k\}$, a *transversal* of $\bigcup_{i \in I} X_i$ is a set B such that $|B| = |I|$ and $|B \cap X_i| = 1$ for each $i \in I$. Let \mathcal{A} consist of all sets A of the form $X_i \cup B$ where $0 \leq i \leq k$ and B is a transversal of $\bigcup_{j \neq i} X_j$. Thus, $|\mathcal{A}| = (k+1)k^k > (n/2)^{n/2}$.

Notice that we may permute the index set of the disjoint union $X = \bigsqcup_{i=0}^k X_i$ and we may independently permute the elements of each X_i without changing the structure of \mathcal{A} . We thus have the following lemma.

Lemma 3.21. *$\text{Aut}(\mathcal{A}) \geq S_{k+1} \times (S_k)^{k+1}$ where S_i is the symmetric group on i letters.* \square

Claim 3.22. *Suppose $k \geq 4$, $0 \leq a_0 \leq a_1 \leq \dots \leq a_k \leq k$ are integers, and $\sum_{i=0}^k a_i \leq 2k$. Then exactly one of the following holds:*

1. $\forall i (a_i + a_{k-i} \leq k)$.
2. $a_0 = a_1 = \dots = a_{k-1} = 1$ and $a_k = k$.

Proof. If the first condition fails, for some $i \leq k/2$ we must have $a_i + a_{k-i} > k$. First we show that $i = 0$: If $i = 1$, then $a_1 + a_{k-1} > k$ and also $a_2 + a_k > k$, since the a_j are increasing. Since $2 < k-1$, this is a contradiction. Similarly, if $i \geq 2$, then $k-1 > k-i$ and $a_k + a_{k-1} > k$, again a contradiction.

Now we show that $a_k = k$: Otherwise, say $a_k = k - j$. Then $a_0 \geq j + 1$. Then $\sum a_i \geq (j + 1)k + (k - i) \geq 2k + (k - i) > 2k$, a contradiction.

Since $a_0 = 1 \leq a_i \leq a_k = k$ for all i and $\sum a_j \leq 2k$, it now follows that $a_1 = \dots = a_{k-1} = 1$ as well, and we are done. \square

To show that \mathcal{A} is minimal, it suffices to verify that the condition of Lemma 3.19 holds. But this follows from Lemma 3.21 and Claim 3.22, letting (after a renumbering if necessary) $a_i = |A \cap X_i|$ for $A \in \mathcal{B}$ for \mathcal{B} a putative intersecting family of $\leq n$ -element sets definable from \mathcal{A} . The point is that if the a_i satisfy conclusion (1) of Claim 3.22, we can let σ be the following involution in S_X : σ exchanges setwise the blocks X_i and X_{k-i} for each i , and in such a way that $A \cap X_i$ is disjoint from $\sigma \cdot (A \cap X_{k-i})$, which is possible since $a_i + a_{k-i} \leq k = |X_i|$. Clearly, $\sigma \in \text{Aut}(\mathcal{A})$, and $A \cap \sigma \cdot A = \emptyset$, so \mathcal{B} was not intersecting after all (recall that $\mathcal{B} \supseteq \mathcal{B}_{\mathcal{A}}$, where $\mathcal{B}_{\mathcal{A}}$ is as in Theorem 3.18). Thus, the a_i satisfy conclusion (2) of Claim 3.22, so $A \in \mathcal{A}$ and since $\text{Aut}(\mathcal{A})$ is transitive on \mathcal{A} , $\mathcal{B} \supseteq \mathcal{A}$.

To prove the general version, consider now the collection \mathcal{A} of sets A as before, but with each block X_i of size pn and a total of $1 + (1 - p)n$ many blocks. A modified version of the claim holds with essentially the same proof, and this gives the result as well. For the sake of exposition, we state a slightly weaker version of the modified statement: Consider blocks of size $n - k$ (for some fixed k) and a total of $k + 1$ blocks. Now we require in the claim that $0 \leq a_0 \leq a_1 \leq \dots \leq a_k \leq n - k$ and $\sum_i a_i \leq n$. The conclusion is that for almost all values of n , either $a_i + a_{k-i} \leq n - k$ for all i , or else $a_0 = \dots = a_{k-1} = 1$ and $a_k = n - k$. \square

Corollary 3.23. *For all $n > 3$, $\psi(2n) > \psi(n)$.* \square

Remark 3.24. Notice that the examples from Theorem 3.20 are low. Also, Corollary 3.23 holds for all n since $\psi(1) = 1$, $\psi(2) = 3$, $\psi(4) \geq \binom{7}{4} = 35$, and $\psi(3) = 10$ by Theorem 3.30 below.

Notice that if $\mathcal{A} \subseteq [X]^n$ is intersecting, then so is

$$\mathcal{A}_+ = \{B \in [X]^{n+1} : \exists A \in \mathcal{A} (A \subseteq B)\}.$$

If \mathcal{A} is minimal and $\mathcal{B} \subseteq [X]^{\leq n}$ is intersecting and definable from \mathcal{A}_+ , then $|\mathcal{B}| \geq |\mathcal{A}|$, since \mathcal{A}_+ is itself definable from \mathcal{A} . Hence, whether \mathcal{A}_+ is $(n + 1)$ -minimal reduces to whether one can define from it a “small” intersecting subfamily of $[X]^{n+1}$.

Notice that $|\mathcal{A}_+| < |\mathcal{A}|$ is possible. For example, let us introduce the notation $[k]^n$ to denote the collection of n -element subsets of $\{0, 1, \dots, k - 1\}$. If $\mathcal{A} = [3]^2$, then $\mathcal{A}_+ = [3]^3$, and if $\mathcal{A} = [5]^3$, then $\mathcal{A}_+ = [5]^4$.

Claim 3.25. *If n is sufficiently large ($n > 34$ suffices) and $\mathcal{A} \subseteq [X]^n$ is n -large, then $|\mathcal{A}_+| > |\mathcal{A}|$.*

Proof. If n is even, by Theorem 3.20, $|\mathcal{A}| \geq (n/2)^{n/2} > 4^n > \binom{2n+1}{n}$, so $|\bigcup A| > 2n + 1$. If n is odd, a similar construction to that the one described in Theorem

3.20 gives the same result: Now consider $(n+1)/2$ blocks, each of size $(n-1)/2$, and consider the family \mathcal{C} of sets that contain one of the blocks and meet the others in exactly one point. This family \mathcal{C} is n -minimal as in Theorem 3.20 and

$$|\mathcal{C}| = \binom{n+1}{2} \left(\frac{n-1}{2} \right)^{\frac{n-1}{2}}.$$

Consider now the set $C = \{(A, B) \in \mathcal{A} \times \mathcal{A}_+ : A \subseteq B\}$. The result follows from a double counting argument: since $|\bigcup A| > 2n+1$, each A in \mathcal{A} can be extended to a B in \mathcal{A}_+ in more than $n+1$ ways, so $|C| > (n+1)|\mathcal{A}|$. On the other hand, each $B \in \mathcal{A}_+$ contains at most $n+1$ members of \mathcal{A} , so $|C| \leq (n+1)|\mathcal{A}_+|$. \square

If one can show that n -largeness of \mathcal{A} implies that \mathcal{A}_+ is $(n+1)$ -minimal, then it follows that $\psi(n) < \psi(n+1)$. Some assumption on \mathcal{A} is necessary, though. For example, if \mathcal{A} is the Fano plane, then \mathcal{A}_+ has size 28 and its complement in $[7]^4$ is also intersecting and has size 7, so \mathcal{A}_+ is not 4-minimal. To establish minimality of \mathcal{A}_+ , it seems that a better understanding of $\text{Aut}(A)$ is required.

Remark 3.26. Notice that $\text{Aut}(\mathcal{A}) \leq \text{Aut}(\mathcal{A}_+)$ since \mathcal{A}_+ is definable from \mathcal{A} .

Question 3.27. If \mathcal{A} is low, does it follow that \mathcal{A}_+ is also low?

3.3 Minimal families of triples

We compile lists of n -minimal families for $n \leq 3$. Let \mathcal{B}_n denote the collection of n -minimal intersecting families of sets, considered up to isomorphism. Then \mathcal{B}_n forms a basis for intersecting families of $(\leq n)$ -element sets in the sense that given any intersecting $\mathcal{A} \subseteq [X]^{\leq n}$ there exists an isomorphic copy of some $\mathcal{A}' \in \mathcal{B}_n$ which is definable from \mathcal{A} .

Recall that $[k]^n$ denotes the set of n -element subsets of $\{0, 1, \dots, k-1\}$. Given a set of integers $A \subseteq \{0, 1, \dots, n-1\}$, let A_n denote the family of translations of A taken modulo n so, for example, $\{013\}_7$ is the Fano plane.

Theorem 3.28. Let L be the family from Example 3.5 and

$$\text{AOct} = \{012, 045, 135, 234\} \subseteq [6]^3.$$

Then:

1. $\mathcal{B}_1 = \{[1]^1\}$,
2. $\mathcal{B}_2 = \{[1]^1, [2]^2, [3]^2\}$,
3. $\mathcal{B}_3 = \{[1]^1, [2]^2, [3]^3, [4]^3, \text{AOct}, \{013\}_5, \{013\}_6, \{013\}_7, [5]^3, L\}$.

The family AOct can be viewed as the collection of red faces of an octahedron whose faces are colored red and blue in such a way that no two adjacent faces share the same color; we call it the *alternating octahedron*.

It is not hard to see that \mathcal{B}_1 and \mathcal{B}_2 are as listed. Furthermore, using the equivalent condition in Theorem 3.18, it is routine to verify that the families in the lists above are minimal. This subsection is devoted to the proof of Theorem 3.28. We proceed by stages: In Theorem 3.30 we show that $\psi(3) = 10$, in Proposition 3.40 we show that the only large members of \mathcal{B}_3 are $[5]^3$ and L , and in Proposition 3.43 we show that no member of \mathcal{B}_3 can have size 8 or 9. Along the way, we also show in Proposition 3.41 that $\xi(1, 3) = 7$; this is an immediate consequence of Theorem 3.28, but providing a direct argument at that point is shorter. We then analyze the remaining possible sizes of members of \mathcal{B}_3 to conclude the proof.

Remark 3.29. Notice that every minimal family in \mathcal{B}_1 , \mathcal{B}_2 , and \mathcal{B}_3 is low. In fact, every minimal family we mention in this paper is also low, suggesting that this might be a general phenomenon.

Theorem 3.30. $\psi(3) = 10$.

Although our argument is specific to the case $n = 3$, we try to illustrate some of the complexities that are present in the analysis of a general large family \mathcal{A} of n -element sets.

Proof. Assume $\mathcal{A} \subseteq [X]^{\leq 3}$ is minimal and $|\mathcal{A}| \geq 10$. Since $\psi(2) = 3$, it then follows that $\mathcal{A} \subseteq [X]^3$ and also that neither S_1 nor S_2 is intersecting. Thus, by Lemma 3.14 we know $d_2 \leq 3$ and $d_1 \leq 3d_2$.

Claim 3.31. $d_1 \geq 4$. Hence, $d_2 \geq 2$.

Proof. This follows from the pigeonhole principle and only requires that $|\mathcal{A}| \geq 8$: Given any $A \in \mathcal{A}$, at least one of its elements belongs to at least three other members of \mathcal{A} . \square

Lemma 3.32. \mathcal{A} is regular.

Proof. Recall that $m_1 = |A \cap S_1|$ for any $A \in \mathcal{A}$. By minimality, \mathcal{A} is regular if and only if $m_1 = 3$. Towards a contradiction, assume \mathcal{A} is not regular, so $m_1 = 2$ by Lemma 3.10. Define d'_2 as

$$d'_2 := \max\{\deg_{\mathcal{A}}(\{x, y\}) : \{x, y\} \in [S_1]^2\}.$$

By minimality of \mathcal{A} , for each $A \in \mathcal{A}$, $A \cap S_1$ is contained in d'_2 elements of \mathcal{A} . Define a graph on S_1 by: $x_1 G x_2$ iff $\deg_{\mathcal{A}}(\{x_1, x_2\}) = d'_2$. Clearly, $d'_2 \leq d_1$.

Claim 3.33. $d'_2 = 1$.

Proof. Assume otherwise. Suppose a, b, c, d are distinct points in S_1 and $ab, cd \in G$. Then there are $x, y \notin S_1$ such that $abx, aby \in \mathcal{A}$, since $d'_2 > 1$. If $\{c, d\} \subset A \in \mathcal{A}$, then A meets at most one of these two sets, contradicting that \mathcal{A} is intersecting.

It follows that $G \subseteq [S_1]^2$ is intersecting. Since $\psi(2) = 3$, from G we can further define an intersecting family of size at most 3 of 2-element sets (or a

singleton). Since this family is definable from \mathcal{A} (because G is), this contradicts that \mathcal{A} is minimal. \square

Our next goal is to show that (S_1, G) is complete.

Claim 3.34. *In (S_1, G) , given any 3 points, there is at most one edge missing. In particular, G is connected of diameter at most 2.*

Proof. Let x, y, z be distinct points in S_1 and suppose x is G -connected to neither y nor z .

Case 1. Suppose first that yGz , i.e., there is $A \in \mathcal{A}$ with $\{y, z\} \subset A$, and let w be the third member of A . Let B be any of the d_1 sets in \mathcal{A} with $x \in B$ and notice that $\{w\} = B \cap A$. It follows that $\deg_{\mathcal{A}}(w) \geq d_1 + 1$, a contradiction.

Case 2. Otherwise. Now let $A \in \mathcal{A}$ be any set containing y , so A contains neither x nor z and any set in \mathcal{A} containing either of them meets A in a point other than y . By the pigeonhole principle at least one element of A must have degree at least $(2d_1 + 1)/2 > d_1$, again a contradiction. \square

Claim 3.35. *(S_1, G) is complete.*

Proof. Suppose otherwise, and let yz be a missing edge. For any $x \in S_1 \setminus \{y, z\}$, both xy and xz are in G . Since $\deg_{\mathcal{A}}(y) = d_1$, there are precisely d_1 such possible vertices x and $|S_1| = d_1 + 2$. Let $A \in \mathcal{A}$ with $\{x, y\} \subset A$ and let a be the third element of A , so $a \notin S_1$. Then for any of the $d_1 - 1$ remaining vertices w of S_1 , if $\{w, z\} \subset B \in \mathcal{A}$ then a is also the third element of B . It follows that $\deg_{\mathcal{A}}(a) \geq d_1$ so $a \in S_1$, a contradiction. \square

We are almost done now: (S_1, G) is complete and $|S_1| = d_1 + 1$. Let $x, y \in S_1$ and let $a \notin S_1$ be such that $\{a, x, y\} \in \mathcal{A}$. Then a also belongs to any set in \mathcal{A} containing 2 of the remaining $d_1 - 1$ elements of S_1 . Hence,

$$\deg_{\mathcal{A}}(a) \geq \binom{d_1 - 1}{2} + 1 \geq d_1$$

since $d_1^2 - 5d_1 + 4 = (d_1 - 4)(d_1 - 1) \geq 0$ as $d_1 \geq 4$, a contradiction. \square

Remark 3.36. For future reference, we explain how to weaken the assumption that $|\mathcal{A}| \geq 10$ in the proof of Lemma 3.32 to $|\mathcal{A}| \geq 8$. The only place where the assumption was used was to conclude that $m_1 \geq 2$ via Lemma 3.10. So, assume that $|\mathcal{A}| = 8$ or 9 and that $m_1 = 1$, thus any $A \in \mathcal{A}$ contains exactly one element of degree d_1 . Clearly, $|S_1| > 1$, by minimality of \mathcal{A} . Arguing as in the proof of Lemma 3.10, if $|S_1| = k + 1$ and $x \in S_1$, then there are kd_1 sets in \mathcal{A} not containing x . Let $A \in \mathcal{A}$ extend $\{x\}$. Those kd_1 sets meet $A \setminus \{x\}$ so one of its two points, say a , satisfies $\deg_{\mathcal{A}}(a) \geq kd_1/2$. Thus $kd_1/2 < d_1$ or $k < 2$, so $k = 1$ and $|S_1| = 2$ so, since $d_1|S_1| = |\mathcal{A}|$, we have that $|\mathcal{A}| = 8$ and $d_1 = 4$.

Let $S_1 = \{z, w\}$ and let $\{x, y, z\}$ extend $\{w\}$, so x or y meets at least two of the four sets extending $\{z\}$, and it follows that $d_{1,2} \geq 3$. Since $d_{1,2} < d_{1,1} = d_1$,

then $d_{1,2} = 3$. Since $3|S_{1,2}|/m_2 = |\mathcal{A}| = 8$ by Fact 3.11, we must have $m_2 \geq 3$. But $m_2 \leq 2$, a contradiction.

Lemma 3.32 means that $X = S_1$, i.e., every point has degree d_1 . By a standard double counting argument, we have $3|\mathcal{A}| = d_1|X|$. We use this fact in conjunction with a more refined version of an earlier counting argument to bound the size of X .

Claim 3.37. $|X| < 8$.

Proof. By the principle of inclusion/exclusion, we have

$$|\mathcal{A}| = \sum_{D \in [A]^1} \deg_{\mathcal{A}}(D) - \sum_{D \in [A]^2} \deg_{\mathcal{A}}(D) + \sum_{D \in [A]^3} \deg_{\mathcal{A}}(D)$$

for any $A \in \mathcal{A}$. Since \mathcal{A} is a regular family of triples, we have

$$|\mathcal{A}| \leq 3d_1 - (d_2 + 2) + 1.$$

Now, since $3d_2 \geq d_1$ and $3|\mathcal{A}| = d_1|X|$, we conclude

$$(|X| - 8)d_1 \leq -3,$$

which implies that $|X| < 8$. \square

Using a similar technique, we can show that \mathcal{A} is not regular on pairs whenever $|X| > 6$. Define m'_2 as

$$m'_2 = |\{D \in [A]^2 : \deg_{\mathcal{A}}(D) = d_2\}|,$$

for any $A \in \mathcal{A}$ (minimality of \mathcal{A} ensures that m_2 is independent of the choice of A).

Claim 3.38. *Suppose that $|X| > 6$. Then $m'_2 < 3$.*

Proof. Suppose, towards a contradiction, that $m'_2 = 3$. Again using

$$|\mathcal{A}| = \sum_{D \in [A]^1} \deg_{\mathcal{A}}(D) - \sum_{D \in [A]^2} \deg_{\mathcal{A}}(D) + \sum_{D \in [A]^3} \deg_{\mathcal{A}}(D),$$

we see

$$|\mathcal{A}| = 3d_1 - 3d_2 + 1.$$

As before, since $3d_2 \geq d_1$ and $3|\mathcal{A}| = d_1|X|$, we may manipulate this to yield

$$(|X| - 6)d_1 \leq 1,$$

which contradicts Claim 3.31 whenever $|X| > 6$. \square

We are now in a position to slightly improve our bound on the size of X .

Claim 3.39. $|X| \neq 7$.

Proof. Suppose towards a contradiction that $|X| = 7$. Since $3|\mathcal{A}| = d_1|X|$, we must have that $|\mathcal{A}|$ is a multiple of 7. By Proposition 2.5, we have $|\mathcal{A}| \leq 25$, so $|\mathcal{A}| \in \{14, 21\}$.

Suppose first that $|\mathcal{A}| = 21$. Since $\binom{7}{2} = 21$ and each element of \mathcal{A} contains three pairs, the average degree of a pair is 3. Since $d_2 \leq 3$, this implies that every pair has degree 3, contradicting Claim 3.38.

We then have that $|\mathcal{A}| = 14$. Mimicking the counting done above, we see that the average degree of a pair is 2. To avoid reaching a contradiction with Claim 3.38, we must then have $d_2 = 3$. Let us now attempt to count the number of pairs of degree 3. Since each $A \in \mathcal{A}$ contains m'_2 pairs of degree 3, $m'_2|\mathcal{A}|$ counts each such pair three times. Thus, there must be $28/3$ pairs of degree 3, which is absurd. \square

Thus, we have $|X| \leq 6$. If $|X| \leq 5$, then certainly $|\mathcal{A}| \leq \binom{5}{3} = 10$, so we may assume $|X| = 6$. Then for any $A \in [X]^3$, at most one of $A, X \setminus A$ is in \mathcal{A} , so $|\mathcal{A}| \leq \binom{6}{3}/2 = 10$. \square

We organized the argument above in a way that allows us to characterize the large $\mathcal{A} \subseteq [X]^{\leq 3}$.

Proposition 3.40. *Let $\mathcal{A} \subseteq [X]^{\leq 3}$ be large. Then either \mathcal{A} is the family of triples from a 5-element set, or \mathcal{A} is isomorphic to the family L of Example 3.5.*

Proof. Using notation as above, we see from the argument of Theorem 3.30 that if \mathcal{A} is a large family of 3-element subsets of X , then $|X| \leq 6$ and if $|X| < 6$, then $|X| = 5$ and $\mathcal{A} = [X]^3$.

Assume now that $|X| = 6$. Then \mathcal{A} is low, by Lemma 3.7. Recall that an intersecting family $\mathcal{B} \subseteq \mathcal{P}(Y)$ is called *maximal* iff for any \mathcal{C} , if $\mathcal{B} \subseteq \mathcal{C} \subseteq \mathcal{P}(Y)$, either \mathcal{C} is no longer intersecting, or else $\mathcal{C} = \mathcal{B}$. Following Meyerowitz [5], we say that an element B of \mathcal{B} is *minimal* iff there is no $A \subset Y$ such that $A \subsetneq B$ and $A \in \mathcal{B}$. It is trivial that any intersecting family is contained in a maximal one and that, if Y is finite, any maximal family has size $2^{|Y|-1}$. In particular, the collection of supersets of sets in \mathcal{A} is maximal and the members of \mathcal{A} are the minimal elements of this maximal family.

In Meyerowitz [5, Proposition 3.1], a list of all 30 maximal intersecting families (up to isomorphism) of a set of size 6 is presented; the families are generated by stages starting from the family of supersets of a singleton $\{a\}$ by means of shifts, and are enumerated according to the number of shifts required to generate them. The families listed in Meyerowitz [5] as $A, B, C, D, E1, E2, F1, F2, G1-G3, H1-H4, I1-I6, J2-J4$ and $K1$ are not the families of supersets of the members of a large minimal \mathcal{A} (in our sense) with $X = \bigcup \mathcal{A}$, since they all have at least one minimal element (in the sense of Meyerowitz [5]) of size other than 3. The families $J1, K2$ and $K3$ are not as required either, since from each of them an element of X is definable. The triple $\{a, b, c\}$ is definable in $K4$. The only remaining example is L , which means that \mathcal{A} must be isomorphic to the family of Example 3.5, which is itself isomorphic to L (and provides a new explanation as to why all the families from Example 3.6 are isomorphic to the one from Example 3.5 as well). \square

In §2 we stated that $\xi(1, 3) = 7$. Theorem 3.30 provides us with an easy way to show this.

Proposition 3.41. $\xi(1, 3) = 7$.

Proof. $\xi(1, 3) \geq 7$ by Theorem 2.10.2. Assume towards a contradiction that $\mathcal{A} \subseteq [X]^3$ is intersecting and such that any non-empty subset of X definable from \mathcal{A} has size at least 8. By passing to a smaller definable intersecting family if necessary, we may assume that \mathcal{A} is minimal, so $|\mathcal{A}| \leq 10$. We use notation as in Fact 3.11 and Theorem 3.30.

Claim 3.42. \mathcal{A} is regular.

Proof. Assume otherwise.

Case 1. There are three distinct degrees of elements of X . Then

$$\begin{aligned} 3|\mathcal{A}| &= d_{1,1}|S_{1,1}| + d_{1,2}|S_{1,2}| + d_{1,3}|S_{1,3}| \geq 8(d_{1,1} + d_{1,2} + d_{1,3}) \\ &\geq 8(3 + 2 + 1) > 3 \times 10 \geq 3|\mathcal{A}|, \end{aligned}$$

a contradiction.

Case 2. There are two distinct degrees. Then

$$3|\mathcal{A}| = d_{1,1}|S_{1,1}| + d_{1,2}|S_{1,2}| \geq 8(2 + 1),$$

so $|\mathcal{A}| \geq 8$, so $d_{1,1} \geq 4$ by Claim 3.31, so

$$3|\mathcal{A}| \geq 8(4 + 1) > 3 \times 10,$$

again a contradiction. □

Then

$$8d_1 \leq |S_1|d_1 = 3|\mathcal{A}| \leq 30,$$

so $d_1 \leq 3$. On the other hand, by Proposition 2.6,

$$8 \leq |S_1| \leq 9 - \frac{6}{d_1},$$

or $6 \leq d_1$, a contradiction. □

Proposition 3.43. *There is no minimal $\mathcal{A} \subseteq [X]^3$ with $|\mathcal{A}| = 8$ or 9.*

Proof. Suppose towards a contradiction that we have a minimal $\mathcal{A} \subseteq [X]^3$ with $|\mathcal{A}| = 8$ or 9. Examining the proof of Theorem 3.30, we only used that $|\mathcal{A}| \geq 8$ to conclude that $d_1 \geq 4$, \mathcal{A} is regular, and $|X| \leq 6$. Thus $|X| = 5$ or 6. A cursory examination of the equation

$$3|\mathcal{A}| = d_1|X|$$

reveals that the only solution fitting the above constraints is $|\mathcal{A}| = 8$, $|X| = 6$, and $d_1 = 4$. Using, once again, the inclusion/exclusion counting as above, we have for any $A \in \mathcal{A}$,

$$|\mathcal{A}| = 3 \times 4 - \sum_{D \in [A]^2} \deg_{\mathcal{A}}(D) + 1,$$

which implies that $\sum_{D \in [A]^2} \deg_{\mathcal{A}}(D) = 5$. In particular, every $A \in \mathcal{A}$ contains a pair D with $\deg_{\mathcal{A}}(D) = 1$; there must therefore be at least 8 such pairs. Thus, at most $\binom{6}{2} - 8 = 7$ pairs have degree greater than 1. Then

$$\begin{aligned} 24 &= 3|\mathcal{A}| \\ &= \sum_{D \in [X]^2} \deg_{\mathcal{A}}(D) \\ &\leq 8 + 7d_2, \end{aligned}$$

so $d_2 = 3$. Since $\sum_{D \in [A]^2} \deg_{\mathcal{A}}(D) = 5$, this means every $A \in \mathcal{A}$ contains one pair of degree 3 and two of degree 1. Thus, there must be a total of 16 pairs of degree 1, a contradiction. \square

Remark 3.44. Let X be the set of vertices of an octahedron \mathcal{H} and let $\mathcal{A} \subseteq [X]^3$ consist of the sets of vertices of the faces of \mathcal{H} . Then $|X| = 6$, $|\mathcal{A}| = 8$, $d_1 = 4$, and \mathcal{A} is low. The family \mathcal{A} is not intersecting, although it is 3-cc. This example may help explain why some care was required in the proof of Proposition 3.43.

We now conclude the proof of Theorem 3.28:

Proof. Suppose that $\mathcal{A} \subseteq [X]^{\leq 3}$ is minimal, and for convenience suppose that $X = \bigcup \mathcal{A}$. Obviously, if $|\mathcal{A}| = 1$ then $\mathcal{A} \in \{[1]^1, [2]^2, [3]^3\}$ and those families are certainly minimal. We thus assume that $|\mathcal{A}| > 1$. In this case $\mathcal{A} \subseteq [X]^3$: if \mathcal{A} contains a set of size 2 or smaller, then we may define from \mathcal{A} one of the elements of \mathcal{B}_2 above, and since $[3]^3$ is definable from $[3]^2$ we can in fact define from \mathcal{A} a family of cardinality 1.

In turn, if the set $Y = \{x \in X : \deg_{\mathcal{A}}(x) = 1\}$ is non-empty, then the family $\mathcal{A}' = \{A \cap (X \setminus Y) : A \in \mathcal{A}\}$ is intersecting, definable from \mathcal{A} , and contains sets of cardinality less than 3. Thus, as above, a family of size 1 is then definable from \mathcal{A}' , and hence from \mathcal{A} . Consequently, we may assume that every point has degree at least 2. Then we have

$$3|\mathcal{A}| = \sum_{x \in X} \deg_{\mathcal{A}}(x) \geq 2|X|,$$

giving the crude bound $|X| \leq \frac{3}{2}|\mathcal{A}|$.

We now consider several cases based upon the value of $|\mathcal{A}|$, recalling that we have already handled $|\mathcal{A}| \geq 8$ and $|\mathcal{A}| = 1$.

$|\mathcal{A}| = 2$: By the above bound, we have $|X| \leq 3$, which prevents this case from being realized.

$|\mathcal{A}| = 3$: We have $|X| \leq 4$. If $|X| < 4$, or if a proper subset of X were definable from \mathcal{A} , then there would be $\mathcal{A}' \subseteq [X]^{\leq 3}$ definable from \mathcal{A} with $|\mathcal{A}'| = 1$, contradicting minimality. Thus, $|X| = 4$ and \mathcal{A} is regular, so we have $3|\mathcal{A}| = d_1|X|$, which has no solutions when $|\mathcal{A}| = 3$ and $|X| = 4$.

$|\mathcal{A}| = 4$: We have $|X| \leq 6$. If $|X| < 4$ or if a proper subset of X were definable from \mathcal{A} , then there would be $\mathcal{A}' \subseteq [X]^{\leq 3}$ definable from \mathcal{A} with $|\mathcal{A}'| = 1$, contradicting minimality. Thus $|X| \geq 4$ and \mathcal{A} is regular, so we have $12 = d_1|X|$. This has two solutions: $|X| = 4, d_1 = 3$, which corresponds to $[4]^3$, and $|X| = 6, d_1 = 2$, which corresponds to AOct.

To see this, note first that if $|X| = 4$, then it is clear that \mathcal{A} must be isomorphic to $[4]^3$, so we focus our attention on the case that $|X| = 6$. As in the proof of Theorem 3.30, we count $|\mathcal{A}|$ by inclusion/exclusion. That is, $|\mathcal{A}| = 4 = 3 \times 2 - \sum_{D \in [A]^2} \deg_{\mathcal{A}}(D) + 1$, so $\sum_{D \in [A]^2} \deg_{\mathcal{A}}(D) = 3$. Thus $d_2 = 1$, so any pair of elements of X is contained in at most one element of \mathcal{A} . Without loss of generality, suppose that $012 \in \mathcal{A}$. Some other element of \mathcal{A} contains 0 but not 1 or 2, without loss let us assume it is 045 . Similarly, an element of \mathcal{A} contains 1 but not 0 or 2. It must also intersect 045 , so, reversing the labels of 4 and 5 if necessary, it is 135 . Since $d_1 = 3$, the last element of \mathcal{A} must be 234 , so $\mathcal{A} = \{012, 045, 135, 234\} = \text{AOct}$.

$|\mathcal{A}| = 5$: We have $|X| \leq 7$. If $|X| < 4$ or if a proper subset of X were definable from \mathcal{A} , then there would be $\mathcal{A}' \subseteq [X]^{\leq 3}$ definable from \mathcal{A} with $|\mathcal{A}'| = 1$, contradicting minimality. Thus $|X| \geq 4$ and \mathcal{A} is regular, so we have $15 = d_1|X|$. This has a unique solution: $|X| = 5, d_1 = 3$.

We count $|\mathcal{A}|$ by inclusion/exclusion, so $5 = 3 \times 3 - \sum_{D \in [A]^2} \deg_{\mathcal{A}}(D) + 1$, or $\sum_{D \in [A]^2} \deg_{\mathcal{A}}(D) = 5$. It cannot be the case that each element of \mathcal{A} contains two pairs of degree 1, since all 10 pairs in $[X]^2$ would have degree 1, contradicting the fact that \mathcal{A} counts 15 pairs (including multiplicity). Thus, it must be the case that each element of \mathcal{A} contains one pair of degree 1 and two pairs of degree 2. This means that five of the pairs in $[X]^2$ have degree 1 and five have degree 2. Suppose that 013 is in \mathcal{A} , with 01 having degree 1 and $13, 03$ having degree 2. Some other element of \mathcal{A} must contain 03 ; call it 023 . Similarly, some other element contains 13 , so it is either 134 or 123 . If it were 123 , then no triple can contain 4 and two of $0, 1, 2, 3$. Thus, $134 \in \mathcal{A}$. Then the only triples that can contain the pair 24 are 024 and 124 ; without loss of generality assume it is 024 . After that, $1, 2$, and 4 are left with degree less than 3, so the last element must contain all of them. Therefore, $\mathcal{A} = \{013, 124, 023, 134, 024\} = \{013\}_5$.

$|\mathcal{A}| = 6$: We have $|X| \leq 9$. If $|X| < 4$ or if a proper subset of X were definable from \mathcal{A} , then there would be $\mathcal{A}' \subseteq [X]^{\leq 3}$ definable from \mathcal{A} with $|\mathcal{A}'| = 1$, contradicting minimality. Thus $|X| \geq 4$ and \mathcal{A} is regular, so we have $18 = d_1|X|$. This has two solutions: $|X| = 9, d_1 = 2$ and $|X| = 6, d_1 = 3$. If $|X| = 9$, then counting $|\mathcal{A}|$ by inclusion/exclusion yields $6 = 2 \times 3 - \sum_{D \in [A]^2} \deg_{\mathcal{A}}(D) + 1$, so $\sum_{D \in [A]^2} \deg_{\mathcal{A}}(D) = 1$, which is absurd.

So, $|X| = 6$. Again counting $|\mathcal{A}|$ we have $6 = 3 \times 3 - \sum_{D \in [A]^2} \deg_{\mathcal{A}}(D) + 1$, so we have $\sum_{D \in [A]^2} \deg_{\mathcal{A}}(D) = 4$. This means each $A \in \mathcal{A}$ contains two pairs

of degree 1 and one pair of degree 2. Then a total of twelve pairs in $[X]^2$ have degree 1, so the other three must have degree 2. The three pairs of degree 2 must be disjoint, since their points of intersection would form a definable set of size at most 3. Without loss of generality, let us label the pairs of degree two 03, 14, 25, and assume that 013 is one of the elements of \mathcal{A} containing 03. The pair 04 must be contained in some element of \mathcal{A} ; since it must contain a pair of degree 2 it must be either 034 or 014. However, if it were 014, the pair 01 would have degree greater than 1, so we must have $034 \in \mathcal{A}$. No element of \mathcal{A} containing 14 can contain 0 or 3 since we have already used the pairs 01, 04, 13, 34, so we must have 124 and $145 \in \mathcal{A}$. Continuing in this fashion gives $\mathcal{A} = \{013, 124, 235, 034, 145, 025\} = \{013\}_6$.

$|\mathcal{A}| = 7$: We have $|X| \leq 10$. If a set of size smaller than 5 were definable from \mathcal{A} , then an intersecting family of size at most 4 would be definable, contradicting minimality. Thus, if \mathcal{A} is not regular it could have at most two distinct degrees, say $d_{1,1}$ and $d_{1,2}$, each witnessed by five points of X . But if this were the case, then we would have

$$21 = 3|\mathcal{A}| = \sum_{x \in X} \deg_{\mathcal{A}}(x) = 5d_{1,1} + 5d_{1,2},$$

which has no solutions. Thus, \mathcal{A} is regular, so we have $21 = d_1|X|$, which implies that $|X| = 7$ and $d_1 = 3$.

Counting $|\mathcal{A}|$ by inclusion/exclusion, we see $7 = 3 \times 3 - \sum_{D \in [A]^2} \deg_{\mathcal{A}}(D) + 1$, so $\sum_{D \in [A]^2} \deg_{\mathcal{A}}(D) = 3$. This means each $A \in \mathcal{A}$ contains three pairs of degree one. Since a total of $3|\mathcal{A}| = 21$ pairs are contained in elements of \mathcal{A} , we conclude that every pair in $[X]^2$ is contained in exactly one element of \mathcal{A} . Without loss of generality, suppose that 013, 026, and 045 are the three elements of \mathcal{A} containing the point zero. Some element of \mathcal{A} contains the pair 12; ruling out the edges already used, the only possibilities are 124 and 125. Switching the labels of 4 and 5 if necessary, we may assume 124 is in \mathcal{A} . Now, some triple must contain 23, and the only possibility is 235. Proceeding in this fashion, the only triple which can contain 46 is 346, and the only triple which can contain 16 is 156. Thus, $\mathcal{A} = \{013, 124, 235, 346, 045, 156, 026\} = \{013\}_7$. \square

4 Low intersecting families

In the previous section we started from an intersecting family $\mathcal{A} \subseteq [X]^{\leq n}$ and investigated bounds on the sizes of intersecting families $\mathcal{A}' \subseteq [X]^{\leq n}$ definable from \mathcal{A} . Here we restrict our attention to those \mathcal{A}' that are themselves subsets of \mathcal{A} .

Definition 4.1. An intersecting family \mathcal{A} is *quasi-low* (*qlow*) iff $\text{Aut}(\mathcal{A})$ acts transitively on \mathcal{A} .

It would seem reasonable to study qlow families $\mathcal{A} \subseteq [X]^n$, since no proper subfamily of such an \mathcal{A} is definable. However, if one is interested in bounding

the size of \mathcal{A} , this is not the right notion to investigate, since no such bound exists:

Example 4.2. Given a set X , natural numbers $m < n$, and $B_0 \in [X]^m$, the family $\mathcal{A} = \{A \in [X]^n : B_0 \subseteq A\}$ is qlow.

Example 4.3. In Example 2.1, assume $|A| = N$. Then \mathcal{A} is qlow of size $3N$.

Hence, we restrict our attention to low families.

Definition 4.4. Let $\rho(n) = \max\{|\mathcal{A}| : \mathcal{A} \subseteq [X]^n \text{ is low and intersecting}\}$.

From the previous section, $\binom{2n-1}{n} \leq \rho(n)$, so $\rho(3) \geq 10$. Similarly, $\rho(n) > (n/2)^{n/2}$ for all $n > 6$. Notice that ρ is well-defined:

Lemma 4.5. *Let $\mathcal{A} \subseteq [X]^n$ be low. If $n = 2$, then $|\mathcal{A}| \leq 3$. If $n \geq 3$, then $|\mathcal{A}| < \binom{n^2-2}{n-1}$.*

Proof. Let $\mathcal{A} \subseteq [X]^n$. If $|X| < 2n$, clearly $|\mathcal{A}| \leq \binom{|X|}{n} \leq \binom{2n-1}{n} \leq \rho(n)$. By Theorem 2.7, $|X| \leq n^2 - 1$. It follows that $\rho(2) = 3$.

If $|X| \geq 2n$ and \mathcal{A} is intersecting, by the Erdős-Ko-Rado theorem, see Erdős-Ko-Rado [3], we have that $|\mathcal{A}| \leq \binom{|X|-1}{n-1}$, so

$$\rho(n) \leq \binom{n^2-2}{n-1}.$$

In fact, if $n \geq 3$ the inequality is strict since $n^2 - 1 > 2n$ and equality in the Erdős-Ko-Rado theorem requires that the family \mathcal{A} consists of all n -element subsets of a set X of size $n^2 - 1$ that contain a fixed element of X . This element is therefore definable from \mathcal{A} , so \mathcal{A} is not low. \square

Theorem 4.6. $\rho(3) = 10$.

Proof. Observe first that to adapt the argument of Theorem 3.30, it suffices to check that a low $\mathcal{A} \subseteq [X]^3$ with $|\mathcal{A}| \geq 10$ satisfies alternative (2) of Lemma 3.14 for $m \in \{1, 2\}$. Aside from these bounds on the degrees, the only use of minimality in the proof of Theorem 3.30 is to ensure that no proper subset of \mathcal{A} is definable from \mathcal{A} ; this holds whenever \mathcal{A} is low.

As usual, we may assume \mathcal{A} is finite. Suppose, towards a contradiction, that either S_1 or S_2 is intersecting. Then, by Theorem 2.7, we may define from \mathcal{A} a subset $Y \subseteq X$ with $|Y| \leq 3$. Since \mathcal{A} is low, we have $|X| \leq 3$, which contradicts $|\mathcal{A}| \geq 10$. \square

We now introduce an operation $*$ that allows us to “lift” small families to larger ones.

Definition 4.7. Let $\mathcal{A} * \mathcal{B} = \{\bigcup_{x \in A} \{x\} \times B_x : B_x \in \mathcal{B}, A \in \mathcal{A}\}$.

Lemma 4.8. *Suppose $\mathcal{A} \subseteq [X]^m$ and $\mathcal{B} \subseteq [Y]^n$ are qlow, and let $\mathcal{C} = \mathcal{A} * \mathcal{B}$. Then $\mathcal{C} \subseteq [X \times Y]^{m \times n}$ is qlow and $|\mathcal{C}| = |\mathcal{A}| |\mathcal{B}|^m$. If \mathcal{A} and \mathcal{B} are low, then so is \mathcal{C} .*

Proof. Notice that \mathcal{C} is intersecting and has the claimed size. Let $C_1, C_2 \in \mathcal{C}$. For $C \in \mathcal{C}$ let $A_C := \text{proj}_1(C)$ be its first coordinate projection, so $A_C = \{x : \exists y (x, y) \in C\}$. There is no loss of generality in assuming that for all $x \in A_{C_1}$, $B_{C_1, x} = B$ is a fixed element of \mathcal{B} , as some automorphism of \mathcal{B} sends $B_{C_1, x}$ to B and this induces an automorphism of \mathcal{C} that sends C_1 to $A_{C_1} \times B$.

Similarly, there is an automorphism of \mathcal{C} sending C_2 to $A_{C_2} \times B$, and there is an automorphism of \mathcal{A} that sends A_{C_1} to A_{C_2} . This automorphism lifts to an automorphism of \mathcal{C} , and appropriately composing these automorphisms we find one that sends C_1 to C_2 , so \mathcal{C} is qlow.

An easy extension of this argument shows that \mathcal{C} is in fact low if \mathcal{A} and \mathcal{B} are low. \square

Definition 4.9. The *powers* of \mathcal{A} are given by $\mathcal{A}^{(1)} = \mathcal{A}$ and $\mathcal{A}^{(k+1)} = \mathcal{A}^{(k)} * \mathcal{A}$.

Fix n and a low $\mathcal{A} \subseteq [X]^n$ such that $|\mathcal{A}| > c^n$, say $|\mathcal{A}| \geq c_1^n$ where $c_1 > c$. Then

$$|\mathcal{A}^{(k)}| = |\mathcal{A}|^{\frac{n^k - 1}{n - 1}},$$

as a straightforward induction establishes, since $|\mathcal{A}^{(k+1)}| = |\mathcal{A}^{(k)}| |\mathcal{A}|^{n^k}$.

If $k > 1$, then $|\mathcal{A}^{(k)}| \geq (c_1^n)^{\frac{n^k - 1}{n - 1}} > c_1^{n^k}$ and $\mathcal{A}^{(k)} \subseteq [X]^{n^k}$. This gives us an infinite sequence of low families that “grow faster” than c^n , obtained by means of our lifting operation.

Remark 4.10. Notice that $*$ is not commutative, but it is associative (up to isomorphism), so $\mathcal{A}^{(k+1)} \cong \mathcal{A} * \mathcal{A}^{(k)}$.

Example 4.11. A “Sierpinski-like” sequence of low families can be obtained by starting with $n = n^1 = 2$ and \mathcal{A} the collection of 2-element subsets of a set of size 3; our construction then produces for infinitely many values of n (namely, the powers of 2) a low \mathcal{A} of n -element subsets, with $|\mathcal{A}| = 3^{n-1}$. In general, we can produce this way, for each k , low families $(\mathcal{A}_n : n \in \mathbb{N})$ with

$$|\mathcal{A}_n| = \left(\binom{2k-1}{k} \right)^{k^{n-1}} \quad \text{and} \quad \mathcal{A}_n \subseteq [X]^{k^n}.$$

Remark 4.12. In general, the operation $*$ does not preserve minimality. To see this, let \mathcal{A} be minimal, let $|Y| = 3$ and consider $\mathcal{B} = \mathcal{A} * [Y]^2$ where each point of $X = \bigcup \mathcal{A}$ is replaced by a “block” of 3 points. By considering invariance under $\text{Aut}(\mathcal{B})$, it follows that the family \mathcal{C} consisting of those $A \in \mathcal{B}$ that contain one of these blocks of size 3 and exactly one point from each other block, is definable from \mathcal{B} , by Lemma A.12, so \mathcal{B} is not minimal. It is worth noting that an analysis of this example led to the lower bounds in Theorem 3.20.

Question 4.13. Is $\text{Aut}(\mathcal{A} * \mathcal{B}) \cong \text{Aut}(\mathcal{A}) * \text{Aut}(\mathcal{B})^n$, where n is the size of the sets in \mathcal{A} ?

Recall that \mathcal{A} is called k -wise t -intersecting iff the intersection of any k members of \mathcal{A} has size at least t , so \mathcal{A} is intersecting iff it is 2-wise 1-intersecting.

It seems worthwhile to explore how the bounds obtained throughout this paper are affected if one now assumes that the family \mathcal{A} is k -wise t -intersecting rather than $(k + 1)$ -cc or simply intersecting.

A Definability

Here we state precisely the notion of definability that we have in mind. We work in the language $\{\in, A\}$ of set theory with one unary predicate symbol. We associate to each set X and family $\mathcal{A} \subseteq \mathcal{P}(X)$ the structure

$$\hat{\mathcal{A}}_X := (\mathcal{P}(X) \sqcup X, \mathcal{A}, \hat{\in}),$$

where \sqcup denotes *disjoint* union, and

$$\alpha \hat{\in} \beta \text{ iff } \alpha \in X, \beta \in \mathcal{P}(X), \text{ and } \alpha \in \beta.$$

We interpret Ax as $x \in \mathcal{A}$ (in particular, Ax implies that $x \in \mathcal{P}(X)$).

The sole purpose of taking disjoint unions and using $\hat{\in}$ rather than \in is to stop the internal set structure of X from introducing unexpected definability; for example, if $X = \{\emptyset, \{\emptyset\}\}$ and $\mathcal{A} = \emptyset$, we could define both elements of X if in $\hat{\mathcal{A}}_X$ we were to use \in rather than $\hat{\in}$, or if we were to take the usual union of X and $\mathcal{P}(X)$, while it is our intention that X should be thought of as a 2-element set with no distinguishing features between its elements. We will gloss over this technicality in what follows.

As usual, a set Y is (*first-order*) *definable from* \mathcal{A} iff there is a first-order formula $\phi(z)$ (where any variable occurring in ϕ other than z is bound) with

$$Y = \{Z : \hat{\mathcal{A}}_X \models \phi(Z)\}.$$

Notice we are not allowing parameters here, since all of our structures are finite and, of course, any finite set is definable with parameters.

Different alternatives to this notion of definability are possible and perhaps some are even more natural. For example, one could argue that rather than first-order definability, the natural notion of definability to consider is that of being *definable in set theory*. For finite families, our simpler setting suffices, as being definable has a natural combinatorial characterization, see Lemma A.12.

Our definitions tend to focus on parameters involving the size of certain sets. In set theory one can define these parameters directly, while in general they are not first-order over $\hat{\mathcal{A}}_X$. However, for any fixed $n \in \mathbb{N}$, we can state that a (definable) set *has precisely n elements*. Since the sizes we consider are finite, we can then overcome the obstacle by a detour: Rather than requiring, for example, that a point x has maximal degree in a graph (this notion not being first-order definable), we can instead require that for a given fixed number k , the point x has degree k . If k happens to be the maximal degree in a specific instance, then this gives (indirectly) the required definition in that instance.

Let S_X denote the *symmetric group* of all permutations of X . Note that the natural action of S_X on X induces an action of S_X on $\mathcal{P}(X)$ given by

$\sigma \cdot A = \sigma[A] := \{\sigma(a) : a \in A\}$, as well as an action of S_X on $\mathcal{P}(\mathcal{P}(X))$ given by $\sigma \cdot \mathcal{A} = \sigma[\mathcal{A}] = \{\sigma \cdot A : A \in \mathcal{A}\}$.

Definition A.1. Given $\mathcal{A} \subseteq \mathcal{P}(X)$, by $\text{Aut}(\mathcal{A})$ we mean $\{\sigma \in S_X : \sigma \cdot \mathcal{A} = \mathcal{A}\}$.

This notation is standard, see for example Meyerowitz [5].

Remark A.2. Using the action mentioned above, we can identify the group of (model theoretic) automorphisms $\text{Aut}(\hat{\mathcal{A}}_X)$ with $\text{Aut}(\mathcal{A})$. Throughout the paper, we use the (slightly imprecise) notation $\text{Aut}(\mathcal{A})$ since X is always clear from context.

Definition A.3. Given a set X and $\mathcal{A} \subseteq \mathcal{P}(X)$, let $Y \subseteq X \cup \mathcal{P}(X)$ and $\sigma \in \text{Aut}(\mathcal{A})$. We say that Y is *invariant* under σ iff $Y = \sigma \cdot Y$ and we say that Y is *invariant* iff it is invariant under all members of $\text{Aut}(\mathcal{A})$.

Proposition A.4. *Given a set X and $\mathcal{A} \subseteq \mathcal{P}(X)$, let $Y \subseteq X \cup \mathcal{P}(X)$. If Y is definable from \mathcal{A} then Y is invariant.* \square

It is thus natural to introduce the following notion:

Definition A.5. Let $\mathcal{A} \subseteq \mathcal{P}(X)$. A set $Y \subseteq X \cup \mathcal{P}(X)$ is *hopelessly undefinable* from \mathcal{A} iff there exists $\sigma \in \text{Aut}(\mathcal{A})$ such that $Y \neq \sigma \cdot Y$.

Remark A.6. If X is finite, the collection $\{Y \subseteq X : Y \text{ is invariant}\}$ is definable from \mathcal{A} . To see this, let $n = |X \cup \mathcal{P}(X)|$ and notice that if $Y \subseteq X$ then Y is invariant iff $\hat{\mathcal{A}}_X \models \phi(Y)$, where $\phi(y)$ is

$$\forall x_1 \dots \forall x_n \forall y_1 \dots \forall y_n \left((\bigwedge_{i < j} x_i \neq x_j \wedge \bigwedge_{i < j} y_i \neq y_j \wedge \bigwedge_i (Ax_i \Leftrightarrow Ay_i) \wedge \bigwedge_{i,j} (x_i \in x_j \Leftrightarrow y_i \in y_j)) \Rightarrow \bigwedge_i (x_i \in y \Leftrightarrow y_i \in y) \right).$$

That is, the formula $\phi(Y)$ states that Y is closed (thus invariant) under any permutation of $X \cup \mathcal{P}(X)$ in $\text{Aut}(\hat{\mathcal{A}}_X) = \text{Aut}(\mathcal{A})$.

Definition A.7. A set $\mathcal{A} \subseteq \mathcal{P}(X)$ is *low* iff every non-empty, proper subset of X is hopelessly undefinable from \mathcal{A} , as is every non-empty, proper subset of \mathcal{A} or, equivalently, iff the group $\text{Aut}(\mathcal{A})$ acts transitively on both X and \mathcal{A} .

In Meyerowitz [5], \mathcal{A} is called *transitive* iff $\text{Aut}(\mathcal{A})$ acts transitively on X . This is a weaker notion than being low.

Example A.8. The following families \mathcal{A} are transitive but not low:

1. $\mathcal{A} = \mathcal{P}(X)$ for any non-empty X .
2. Let X be the set of vertices of a hexagon, labeled clockwise as 1–6, and let \mathcal{A} consist of the rotations of the triangles $\{1, 2, 4\}$ and $\{1, 3, 5\}$.

We are mainly interested in the case where the family \mathcal{A} is intersecting. The following are typical examples of low intersecting families.

Example A.9. Let X be the ordered set of size \aleph_1 defined by considering a set A of order type ω_1 , and replacing each element of A with a copy of \mathbb{Q} . Let \mathcal{A} be the collection of countable non-empty subsets of X closed under predecessors and whose supremum does not exist. Since all countable dense linear orders without endpoints are order isomorphic, it follows that \mathcal{A} is low.

Example A.10. Let X be an infinite set. Let \mathcal{U} be a uniform ultrafilter on X , and let \mathcal{A} be the collection of sets in \mathcal{U} whose complement also has size $|X|$. Then \mathcal{A} is low. (For the easy proof, see Caicedo-Clemens-Conley-Miller [1].)

Example A.11. Let X be a set of size 10 and let \mathcal{A} be the collection of all subsets of X of size 6. Then \mathcal{A} is low.

Additional examples can be found throughout the paper; in particular, see Theorem 2.10.

It turns out that for *finite* structures, a set is definable iff it is invariant under automorphisms:

Lemma A.12. *Assume X is finite and let $B \subseteq \mathcal{P}(X) \cup X$. Then B is definable from \mathcal{A} iff it is invariant under $\text{Aut}(\mathcal{A})$.*

Proof. This is an easy consequence of either Beth's definability theorem or Svenonius's theorem, see Hodges [4, §10.5]. Briefly: Let $\vec{a} = \langle a_1, \dots, a_n \rangle$ enumerate $X \sqcup \mathcal{P}(X)$, where $n = |X| + 2^{|X|}$, and let $\phi_{\mathcal{A}}^{\vec{a}}(\vec{x})$ be the formula

$$\bigwedge_{i < j} x_i \neq x_j \wedge \forall y \bigvee_i y = x_i \wedge \bigwedge_{a_i \in a_j} x_i \in x_j \wedge \bigwedge_{a_i \notin a_j} x_i \notin x_j \wedge \bigwedge_{a_i \in \mathcal{A}} Ax_i \wedge \bigwedge_{a_i \notin \mathcal{A}} \neg Ax_i,$$

where $i, j \in \{1, \dots, n\}$ in all connectives. Suppose $\hat{\mathcal{B}}_Y \models \exists \vec{y} \phi_{\mathcal{A}}^{\vec{a}}(\vec{y})$, as witnessed by \vec{b} . Then the map $b_i \mapsto a_i$ is an isomorphism between $\hat{\mathcal{B}}_Y$ and $\hat{\mathcal{A}}_X$.

Assume now that $Y \subseteq \mathcal{P}(X) \cup X$ is invariant under $\text{Aut}(\mathcal{A})$ and let $\varphi(z)$ be the formula $\exists \vec{x} \psi(\vec{x}, z)$, where $\psi(\vec{x}, z)$ is the formula

$$\phi_{\mathcal{A}}^{\vec{a}}(\vec{x}) \wedge \bigwedge_{a_i \notin Y} z \neq x_i \wedge \bigvee_{a_i \in Y} z = x_i.$$

Then Y is definable from \mathcal{A} via φ : The tuple \vec{a} witnesses that $Y \subseteq \{a : \varphi(a)\}$, and the invariance of Y under automorphisms guarantees that for any tuple \vec{b} , $\{a : \psi(\vec{b}, a)\}$ is either empty (if $\phi_{\mathcal{A}}^{\vec{a}}(\vec{b})$ fails) or Y . \square

Corollary A.13. *If X is finite and $B \subseteq \mathcal{P}(X) \cup X$, then B is not definable from \mathcal{A} iff B is hopelessly undefinable from \mathcal{A} .* \square

References

- [1] A. Caicedo, J. Clemens, C. Conley, and B.D. Miller. *Defining small sets from κ -cc families of subsets. II: Families of countable sets*. Preprint, (2009).

- [2] J. Clemens, C. Conley, and B.D. Miller. *Borel homomorphisms of smooth σ -ideals*. Submitted to the Annals of Pure and Applied Logic, (2007).
- [3] P. Erdős, C. Ko, R. Rado. *Intersection theorems for systems of finite sets*. Quarterly Journal of Mathematics, Oxford Series, series 2. **12** (1961), 313–320.
- [4] W. Hodges. **Model theory**. Cambridge University Press, (1993).
- [5] A. Meyerowitz. *Maximal intersecting families*. European Journal of Combinatorics. **16** (1995), 491–501.
- [6] J.H. van Lint, R.M. Wilson. **A course in combinatorics**. Cambridge University Press, (2001), 2nd edition.

AMS (2000) Subject Classification: Primary 05D05, 03B10, secondary 05B30.
Keywords: Intersecting families, chain conditions, definability.

Andrés Eduardo Caicedo
 Boise State University
 Department of Mathematics
 1910 University Drive
 Boise, ID 83725-1555
Homepage: <http://math.boisestate.edu/~caicedo/>
Email: caicedo@math.boisestate.edu

John D. Clemens
 Penn State University
 Mathematics Department
 University Park, PA 16802
Homepage: <http://www.math.psu.edu/clemens/>
Email: clemens@math.psu.edu

Clinton Conley
 University of California, Los Angeles
 Mathematics Department — Box 951555
 Los Angeles, CA 90095-1555
Homepage: <http://www.math.ucla.edu/~clintonc/>
Email: clintonc@math.ucla.edu

Benjamin Miller
 8159 Constitution Road
 Las Cruces, NM 88007
Homepage: <http://glimmeffros.googlepages.com>
Email: glimmeffros@gmail.com